

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.				
1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE October 1997		3. REPORT TYPE AND DATES COVERED
4. TITLE AND SUBTITLE  Changes in peak oxygen uptake and plasma volume in fit and unfit subjects following exposure to a simulation of microgravity			5. FUNDING NUMBERS  PE - 77 PR- 55 TA- B5 WU- 01	
6. AUTHOR(S)  Victor A. Convertino				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Armstrong Laboratory (AFMC) Aerospace Medicine Directorate Clinical Sciences Division, Physiological Research Branch 2507 Kennedy Circle Brooks AFB, TX 78235-5117			8. PERFORMING ORGANIZATION REPORT NUMBER  AL/AO-JA-1997-0094	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)			10. SPONSORING/MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION/AVAILABILITY STATEMENT  Approved for public release; distribution is unlimited.			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words)  To test the hypothesis that the magnitude of reduction in plasma volume and work capacity following simulated microgravity is dependent on the initial level of aerobic fitness, maximal oxygen uptake (Vo2max) was measured in a group of physically fit subjects and compared to Vo2max in a group of relatively unfit subjects before and after 10 days of continuous 6° headdown tilt (HDT). Ten fit subjects (40 ± 2 yr) with mean ± SE VO2max = 48.5 ± 1.9 ml/kg/min were matched for age, height, and lean body weight with 10 unfit subjects (VO2max = 38.2 ± 1.8 ml/kg/min). Before and after HDT, plasma, blood, and red cell volumes and body composition were measured and all subjects underwent a graded supine cycle ergometer test to determine VO2max. The reduction in VO2max in fit subjects (-16.2%, P < 0.05) was greater (P < 0.05) than that of unfit subjects (-6.1%, P < 0.05). Similarly, reductions in plasma volume (-18.3%, P < 0.05) and blood volume (-16.0%, P < 0.05) in fit subjects were larger (P < 0.05) than those of unfit subjects (blood volume = -5.6%, P < 0.05; plasma volume = -6.6%, P < 0.05). Reduction in plasma volume was associated with greater (P < 0.05) negative body fluid balance during the initial 24 h of HDT in the fit group (912 ± 154 ml) compared to unfit subjects (453 ± 200 ml). Red cell volume was unchanged in the unfit subjects but decreased by 12.2% (P < 0.05) in the fit subjects. Percent change for VO2max correlated with percent change in plasma volume (r = +0.79, P < 0.05). Following exposure to simulated microgravity, fit subjects demonstrated a larger reduction in VO2max than unfit subjects which was associated with larger reductions in plasma and blood volume.				
14. SUBJECT TERMS Bed rest, blood volume, fitness, maximal oxygen uptake, microgravity, fluid intake, urine output			15. NUMBER OF PAGES  20	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT  UL	

# Changes in peak oxygen uptake and plasma volume in fit and unfit subjects following exposure to a simulation of microgravity

V.A. CONVERTINO

Physiology Research Branch, Clinical Sciences Division, Armstrong Laboratory, Brooks AFB, TX, USA

## ABSTRACT

To test the hypothesis that the magnitude of reduction in plasma volume and work capacity following exposure to simulated microgravity is dependent on the initial level of aerobic fitness, peak oxygen uptake ( $\dot{V}O_{2peak}$ ) was measured in a group of physically fit subjects and compared with  $\dot{V}O_{2peak}$  in a group of relatively unfit subjects before and after 10 days of continuous 6 head-down tilt (HDT). Ten fit subjects ( $40 \pm 2$  year) with mean  $\pm$  SE  $\dot{V}O_{2peak} = 48.9 \pm 1.7$  mL kg<sup>-1</sup> min<sup>-1</sup> were matched for age, height, and lean body weight with 10 unfit subjects ( $\dot{V}O_{2peak} = 37.7 \pm 1.6$  mL kg<sup>-1</sup> min<sup>-1</sup>). Before and after HDT, plasma, blood, and red cell volumes and body composition were measured and all subjects underwent a graded supine cycle ergometer test to determine  $\dot{V}O_{2peak}$ . Reduced  $\dot{V}O_{2peak}$  in fit subjects (-16.2%) was greater than that of unfit subjects (-6.1%). Similarly, reductions in plasma (-18.3%) and blood volumes (-16.0%) in fit subjects were larger than those of unfit subjects (blood volume = -5.6%; plasma volume = -6.6%). Reduced plasma volume was associated with greater negative body fluid balance during the initial 24 h of HDT in the fit group ( $912 \pm 154$  mL) compared with unfit subjects ( $453 \pm 200$  mL). Per cent change for  $\dot{V}O_{2peak}$  correlated with percentage change in plasma volume ( $r = +0.79$ ). Following exposure to simulated microgravity, fit subjects demonstrated larger reductions in  $\dot{V}O_{2peak}$  than unfit subjects which was associated with larger reductions in plasma and blood volume. These data suggest that the magnitude of physical deconditioning induced by exposure to microgravity without intervention of countermeasures was influenced by the initial fitness of the subjects.

**Keywords** bedrest, blood volume, fitness, fluid intake, microgravity, peak oxygen uptake, urine output.

Received 13 October 1997, accepted 29 May 1998

The reduction in maximal or peak oxygen uptake ( $\dot{V}O_{2peak}$ ) following exposure to spaceflight (Levine *et al.* 1996) and ground simulations of low gravity (Convertino 1995) is well recognized. However, the contribution of physical fitness to the magnitude of loss in aerobic power induced by microgravity is unclear. The observation that the magnitude of increase in  $\dot{V}O_{2peak}$  with endurance exercise training is limited by pre-training  $\dot{V}O_{2peak}$  (a so-called 'ceiling effect') (Saltin *et al.* 1969) leads to the logical hypothesis that individuals with high  $\dot{V}O_{2peak}$  may experience greater reduction in  $\dot{V}O_{2peak}$  during exposure to reduced physical activity compared with their less aerobically fit counterparts. This hypothesis has important implications for development and application of exercise management

for individuals confined to clinical and low gravity environments. The notion that greater baseline aerobic fitness might be associated with larger reduction in  $\dot{V}O_{2peak}$  was supported by the early observation that a 22% reduction in  $\dot{V}O_{2peak}$  was measured following 28 days of bedrest in a subject with initial  $\dot{V}O_{2peak}$  of 4.15 L min<sup>-1</sup> compared with 13% reduction observed in a subject with initial  $\dot{V}O_{2peak}$  of 3.54 L min<sup>-1</sup> (Taylor *et al.* 1949). This relationship was further corroborated by the observations that four endurance-trained runners who underwent 6 h of water immersion reduced their  $\dot{V}O_{2peak}$  by 19% compared with only 10% in four untrained subjects (Stegemann *et al.* 1969) and that pre-bedrest  $\dot{V}O_{2peak}$  was significantly correlated with percentage reduction in  $\dot{V}O_{2peak}$  in 15 subjects after

Correspondence: Victor A. Convertino PhD, Physiology Research Branch, AL/AOCY, 2507 Kennedy Circle, Brooks AFB, TX 78235-5117, USA.

14 days bedrest (Convertino *et al.* 1986). However, analysis of data from other investigations (Chase *et al.* 1966, Saltin *et al.* 1968) did not support a close relationship between initial  $\dot{V}O_{2peak}$  and the magnitude of reduction in  $\dot{V}O_{2peak}$  during low gravity analogues, a finding which may have been compounded by high experimental variability and low sample sizes (Greenleaf & Kozlowski 1982).

In addition to high  $\dot{V}O_{2peak}$ , cross-sectional comparisons have demonstrated that athletes and fit individuals have larger plasma and blood volumes compared with sedentary populations (Convertino 1991), a relationship that is closely associated with  $\dot{V}O_{2peak}$ . As exposure to low gravity environments and their analogues cause hypovolemia (Fischer *et al.* 1967, Johnson *et al.* 1975, Greenleaf *et al.* 1989, Convertino 1990, 1995, 1996, 1997), it seems reasonable to expect that larger reductions in  $\dot{V}O_{2peak}$  following exposure to microgravity may be related to larger reductions in plasma volume.

The relationship between fitness level and magnitude of reductions in  $\dot{V}O_{2peak}$  and plasma volume has not been systematically examined. Therefore,  $\dot{V}O_{2peak}$ , plasma volume, and body fluid balance were measured in a group of aerobically fit subjects before and after exposure to 10 days of head-down tilt (HDT) and compared with changes in an unfit group to test the hypothesis that greater aerobic capacity would be associated with larger reduction in  $\dot{V}O_{2peak}$  and circulating vascular volume.

## METHODS

### Subjects

Twenty healthy men, with a mean age  $\pm$  SE of  $41 \pm 1$  year (range 35–51), a mean height of  $178 \pm 1$  cm (range 168–185), a mean weight of  $77.2 \pm 1.9$  kg (range 64.0–92.9), and a mean relative body fat of  $19.9 \pm 1.2\%$  (range 9.8–28.9) were selected to participate in this study. After selection, the subjects signed a written consent form indicating that they understood the purpose of the study, nature of all tests, foreseeable inconvenience, discomforts, risks,

restrictions and controls. All experimental procedures and protocols were approved by the Human Research Review Boards of the National Aeronautics and Space Administration (NASA) at Ames Research Centre. Selection of subjects was based on results of a screening evaluation comprising a detailed medical history, physical examination, blood chemistry analysis, urinalysis, chest X-ray, and electrocardiogram. Subjects were familiarized with all laboratory personnel, procedures, protocols, mechanics of cycle ergometry in the supine position, and underwent a graded treadmill test to determine their maximal oxygen uptake ( $\dot{V}O_{2peak}$ ) during a 5-day orientation period conducted prior to the study.

### General protocol

Subjects were divided into fit and unfit groups ( $N = 10$  subjects each) according to the 10 highest and 10 lowest treadmill maximal oxygen uptake ( $\dot{V}O_{2peak}$ ) values. This procedure produced two groups that were matched for age, height, and lean body mass (Table 1). The overall experimental protocol consisted of a 7-day ambulatory baseline period (pre-HDT) followed by 10 days of continuous bedrest in the anti-orthostatic 6 HDT position. During HDT, subjects remained in the head-down position continuously for all activities including excretory functions, showering, and eating. During this 17-day experimental period, subjects were housed 24 h day<sup>-1</sup> in the Human Research Facility at NASA-Ames Research Centre and followed a controlled diet. The average daily caloric intake was 2500–2800 kcal ( $\approx 45\%$  carbohydrate, 38% fat, 17% protein). Dietary sodium and potassium were held constant at  $\approx 120$  and 60–80 mEq day<sup>-1</sup>, respectively. Fluid intake was *ad libitum*. The photoperiod was 16 h light to 8 h dark with lights on at 0700 h. The 6 HDT was chosen because actual changes in cardiovascular responses induced by spaceflight have been closely simulated by this ground-based analogue (Convertino 1995). Each subject underwent measurement of their plasma volume and supine ergometer  $\dot{V}O_{2peak}$  on day 7 of the pre-HDT and on the final day of HDT (post-HDT).

	Unfit ( $N = 10$ )	Fit ( $N = 10$ )	<i>t</i> value	<i>P</i> value
Age, year	$40 \pm 2$	$42 \pm 2$	1.746	0.098
Height, cm	$178 \pm 2$	$178 \pm 2$	0.238	0.815
Weight, kg	$80.9 \pm 2.4$	$73.4 \pm 2.6$	2.135	0.047
Lean mass, kg	$59.1 \pm 2.4$	$59.6 \pm 2.8$	0.145	0.887
Body fat, %	$21.8 \pm 0.8$	$13.8 \pm 1.2$	5.340	<0.0001
Blood volume, mL kg <sup>-1</sup>	$70.4 \pm 1.9$	$79.1 \pm 1.6$	3.472	0.003
$\dot{V}O_{2peak}$ , mL kg <sup>-1</sup> min <sup>-1</sup>	$37.7 \pm 1.6$	$48.9 \pm 1.7$	4.882	<0.0001

Values are mean  $\pm$  1SE

Table 1 Subject descriptive data

### Measurements of maximal oxygen uptake

Each subject performed a maximal exercise tolerance test in the supine position on a Collins electronic cycle ergometer. The exercise test used to determine  $\dot{V}O_{2\text{peak}}$  began with a 4-min warm-up period of no resistance pedalling and increased 15 W each minute until volitional exhaustion. The supine position was chosen to minimize the orthostatic factor which contributes to the reduction in work tolerance following HDT (Convertino 1995). Oxygen uptake and heart rate were measured during each 30 s of the exercise test.

The subjects used an Otis-McKerrow respiratory valve and the volume of expired gas was measured by a Parkinson-Cowan high-velocity, low-resistance meter. A potentiometer at the gas meter dial transmitted electrical outputs to a two-channel recorder (MRE Model M22). In this way, expired ventilation flow rate could be recorded. The expired gas was continuously extracted via a pump (Scientific Products) from a 5-L mixing chamber (R-Pel) which was placed between the subject and the gas meter to a 2-L gas bag (Ohio Medical) attached to a Costill-Wilmore valve. The ventilation flow rate was corrected by adding the pump withdrawal volume rate of  $2.5 \text{ L min}^{-1}$  to the recorded volume of expired air. The composition of expired gas was analysed for the fraction of mixed expired oxygen and carbon dioxide using mass spectrometry (Perkin-Elmer). The gas analysers were calibrated with known oxygen and carbon dioxide concentrations before and after each test.

### Measurements of plasma volume

Plasma volume was determined using a modified Evans blue dye dilution technique (Greenleaf *et al.* 1979, Convertino *et al.* 1996). After each subject was stabilized in the supine position for 30 min, an intravenous injection of 11.5 mg of dye diluted with isotonic saline solution (2.5 mL) was administered. The dye from a 10-min post-injection blood sample, recovered from the plasma with a wood-cellulose powder (Solka-Floc SW-40 A) chromatographic column, was compared with a standard dye solution at 615 nm with a spectrophotometer. Peripheral venous blood for duplicate microhematocrit (Hct) determinations was drawn into capillary tubes and centrifuged for 10 min. Total circulating blood volume (BV) and red cell volume were calculated from the PV and Hct. Total blood volume was calculated from the plasma volume and hematocrit measurements. Using these procedures in our laboratory, test-retest correlation coefficient for blood volume was 0.969 ( $N = 12$ ) and the average changes were 82 mL (average  $\% \Delta = 1.5\%$ ,  $N = 17$ ), 75 mL (average  $\% \Delta = 1.5\%$ ,  $N = 19$ ), and 56 mL (average  $\% \Delta = 1.1\%$ ,  $N = 23$ ) when measurements were determined 4, 8, and 15 days apart, respectively

(Greenleaf *et al.* 1979), and was 0.881 ( $N = 7$ ) with average change of 25 mL (average  $\% \Delta = 0.7\%$ ) when measurements were determined 11 months apart (Convertino *et al.* 1996).

### Assessment of 24-h fluid intake and urine output

Fluid intakes and urine outputs were measured daily for total volume to the nearest 1 mL. During HDT, subjects remained in the head-down posture to urinate into a sterile urinal. All 24-h volumes were calculated from the sum of all volumes from samples collected between the start of the second urine void of each morning through and including the first urine void of the following day. The group mean for 24-h intake and output volumes during pre-HDT was determined from the average volumes collected during the final 6 days of the baseline ambulatory period while a value calculated to represent the mean 24-h intake and output volumes during HDT was determined by averaging individual 24-h values over the days of HDT.

Data were presented as mean  $\pm 1$  standard error ( $\pm \text{SE}$ ). Differences between pre-HDT and post-HDT alterations in  $\dot{V}O_{2\text{peak}}$ , plasma, blood and red cell volumes, fluid intake, urine output, and net body fluid balance were analysed by Student's *t*-test statistics for independent groups.

## RESULTS

### Body composition


Although group matching produced no statistically discernable differences in age, height, and lean body mass, fit subjects were characterized by less fat and body weight and greater blood volume (Table 1). Body weight and lean body mass remained essentially unchanged ( $P = 0.3935$ ) in both the fit ( $0.2 \pm 0.5\%$  and  $-0.08 \pm 0.4\%$ , respectively) and the unfit ( $0.3 \pm 0.4\%$ ) groups after HDT. HDT increased percentage body fat ( $P < 0.05$ ) in fit ( $\% \Delta = 4.2 \pm 1.9$ ) and unfit ( $\% \Delta = 3.8 \pm 1.9$ ) subjects.

### $\dot{V}O_{2\text{peak}}$

According to the experimental design, average baseline  $\dot{V}O_{2\text{peak}}$  was 23% higher in fit than unfit subjects (Table 1). HDT decreased  $\dot{V}O_{2\text{peak}}$  from  $2.87 \pm 0.17$  to  $2.41 \pm 0.13 \text{ L min}^{-1}$  and from  $2.40 \pm 0.15$  to  $2.24 \pm 0.13 \text{ L min}^{-1}$  in the fit and unfit subjects, respectively. The percentage reduction in  $\dot{V}O_{2\text{peak}}$  in the fit group was more than double ( $t = 4.597$ ,  $P = 0.0002$ ) that of the unfit group (Fig. 1).

### Plasma volume

HDT decreased baseline blood volume in both groups primarily as a result of a mean reduction in plasma




**Figure 1** Per cent reductions in peak oxygen uptake ( $\dot{V}O_{2peak}$ ), total blood volume (TBV), red blood cell volume (RBCV), and plasma volume (PV) after 10 days of HDT in unfit (□) and fit (■) subjects. Lines represent  $\pm$  1SE and \* indicate  $P < 0.05$  for fit vs. unfit groups.

volume (Fig. 1). A plasma volume reduction of 18.3% in the fit subjects from  $47.9 \pm 1.5 \text{ mL kg}^{-1}$  before HDT to  $39.1 \pm 1.2 \text{ mL kg}^{-1}$  after HDT was greater ( $t = 6.140$ ,  $P < 0.0001$ ) than the 6.6% reduction measured in the unfit group (from  $43.4 \pm 1.3 \text{ mL kg}^{-1}$  before HDT to  $38.7 \pm 1.1 \text{ mL kg}^{-1}$  after HDT). As a result of plasma volume reductions, blood volume was decreased proportionately more ( $t = 7.074$ ,  $P < 0.0001$ ) in the fit subjects compared with their unfit counterparts (Fig. 1). A reduction in red blood cell volume from  $2268 \pm 100 \text{ mL}$  before HDT to  $2013 \pm 92 \text{ mL}$  after HDT in the fit subjects was greater ( $t = 3.307$ ,  $P = 0.004$ ) than the statistically unchanged RBC volume measured in the unfit group (from  $2214 \pm 83 \text{ mL}$  before HDT to  $2131 \pm 61 \text{ mL}$  after HDT).


#### Fluid intake and urine output

Measurements for average 24-h fluid intakes and urine outputs during the pre-HDT baseline period and 10 days of HDT are presented in Fig. 2. HDT induced a transient diuresis during the initial 24 h of exposure, increasing urine flow from an average 6-day pre-HDT baseline level of  $2070 \pm 230$  to  $2631 \pm 267 \text{ mL}$  after the first day of HDT in the fit group ( $+561 \pm 136 \text{ mL}$ ) that was greater ( $t = 2.208$ ,  $P = 0.040$ ) than the increase in urine output from  $2308 \pm 275$  to  $2456 \pm 274 \text{ mL}$  ( $+148 \pm 128 \text{ mL}$ ) in the unfit group. HDT decreased fluid intake in a similar magnitude ( $t = 0.048$ ,  $P = 0.962$ ) in both groups, from an average 6-day pre-HDT baseline level of  $2107 \pm 226$  to



**Figure 2** Average 24-h urine outputs (a), fluid intakes (b), and net fluid balance (c) before (Baseline) and during 10 days of 6° head-down tilt (6° HDT) in unfit (○, - - -) and fit (●, ····) subjects. Lines represent  $\pm$  1SE.

$1619 \pm 49 \text{ mL}$  during HDT in the fit group ( $-388 \pm 63 \text{ mL}$ ) compared with a decrease from  $2399 \pm 270$  to  $2123 \pm 37 \text{ mL}$  ( $-276 \pm 159 \text{ mL}$ ) in the unfit group. As result of increased urine output and



**Figure 3** Relationship between percentage changes (%Δ) in plasma volume and  $\dot{V}O_{2peak}$  after 10 days of HDT in 10 fit (●) and 10 unfit (○) subjects.

reduced fluid intake, a net negative fluid balance of  $912 \pm 154$  mL induced by the initial 24 h of HDT in the fit group was twice as great ( $t = 1.819$ ,  $P = 0.086$ ) as the net negative fluid balance of  $453 \pm 200$  mL in the unfit group (Fig. 2).

#### Relationship between %Δ $\dot{V}O_{2peak}$ and %Δ plasma volume

Figure 3, generated from the individual data sets of the 10 fit and 10 unfit subjects, illustrates the relationship between percentage changes in plasma volume and  $\dot{V}O_{2peak}$  as a result of HDT. This analysis produced a regression equation where %Δ $\dot{V}O_{2peak} = 0.76$  [%Δ plasma volume] & minus;1.7, with a correlation coefficient of 0.787.

## DISCUSSION

Investigations limited by variabilities from small sample sizes and cross-study comparisons in experimental protocols have provided anecdotal evidence that both support (Taylor *et al.* 1949, Stegemann *et al.* 1969, Convertino *et al.* 1986) or reject (Chase *et al.* 1966, Saltin *et al.* 1968, Greenleaf and Kozlowski 1982) the hypothesis that a larger baseline aerobic fitness is associated with greater reduction in  $\dot{V}O_{2peak}$  following exposure to a ground-based analogue of low gravity. A strength of the present study was the use of an experimental protocol specifically designed to test this hypothesis with a sample size that provided adequate statistical power. The major finding from the present study was consistent with the concept that individuals with higher aerobic capacity have greater reduction in

$\dot{V}O_{2peak}$  during adaptation to microgravity than less fit individuals as the reduction in  $\dot{V}O_{2peak}$  of a group of aerobically fit subjects was more than two times that of a group of aerobically unfit subjects. These results also suggest that levels of physical conditioning may influence the degree of adaptation that occurs with prolonged exposure to microgravity.

The mechanisms of reduction in  $\dot{V}O_{2peak}$  during exposure to low gravity conditions is clearly multifaceted. One of the most consistent and rapid adaptations to microgravity is the reduction in blood volume (Fischer *et al.* 1967, Johnson *et al.* 1975, Convertino 1990, 1995, 1996). The significant correlation of 0.79 between %Δ in plasma volume and %Δ in  $\dot{V}O_{2peak}$  observed in the present study suggests that reduction in circulating vascular volume was a major contributing factor to the magnitude of loss in aerobic capacity in fit and unfit subjects. This coefficient was similar to the correlation of 0.84 between %Δ in plasma volume and %Δ in  $\dot{V}O_{2peak}$  generated from a cross-sectional compilation of data from 12 independent investigations (Convertino 1995). High correlation coefficients between the relative changes in plasma volume and  $\dot{V}O_{2peak}$  suggest that microgravity-induced hypovolemia probably contributes importantly to the reduction in the capacity of the body to transport and utilize oxygen during exercise.

It is well established that the reduction in  $\dot{V}O_{2peak}$  following exposure to actual or ground-based analogues of microgravity is primarily the result of reduced stroke volume (Saltin *et al.* 1968, Levine *et al.* 1996). As high  $\dot{V}O_{2peak}$  is associated with large blood volumes (Convertino 1991) and deconditioning-induced hypovolemia is associated with reduced maximal stroke volume and  $\dot{V}O_{2peak}$  (Coyle *et al.* 1986), it is not surprising that the fit subjects in the present study who demonstrated greater reduction in their  $\dot{V}O_{2peak}$  also had proportionately larger hypovolemia compared with the unfit subjects (Fig. 1). The adverse impact of greater hypovolemia on stroke volume and  $\dot{V}O_{2peak}$  following exposure to microgravity may be further exaggerated by a steeper Frank-Starling curve that has been reported in aerobically fit subjects (Levine 1993). A causal relationship between the magnitude of hypovolemia and reduction in  $\dot{V}O_{2peak}$  is further supported by the observation that reduction in  $\dot{V}O_{2peak}$  after HDT has been minimal when hypovolemia was ameliorated (Greenleaf *et al.* 1989). Although several mechanisms may contribute, it is clear that the greater reduction in  $\dot{V}O_{2peak}$  in fit individuals may be attributed to the association between their larger decreases in blood volume and left ventricular filling.

Reductions in plasma volume during HDT in the present investigation reflected alterations in balance between fluid intake and urine output induced by

changed  
to  
mitch  
list!

exposure to this ground-based analogue of microgravity. Generally, there was a negative fluid balance during the initial 24 h of HDT in all subjects resulting from a diuresis and attenuated fluid intake (Fig. 3). After the initial 24 h of HDT, average fluid intake and urine output were attenuated proportionately throughout the remaining 9 days of HDT, maintaining fluid balance within pre-HDT levels. While the attenuation of fluid intake during HDT was similar between fit and unfit subjects, greater reduction in plasma volume observed in the fit subjects was associated with their nearly fourfold greater diuresis during the initial 24 h of HDT. Greater urine flow in fit compared with unfit subjects in the present study was similar to findings of a previous investigation that demonstrated a 22% greater diuresis in trained runners compared with sedentary subjects during 5 h of head-out water immersion (Convertino *et al.* 1993). Mechanisms underlying greater hypovolemia and diuresis during HDT in fit compared with unfit subjects may include relatively larger central blood volume shifts associated with greater baseline blood volume in aerobically fit individuals, resulting in greater glomerular filtration and inhibition of anti-diuretic hormone and aldosterone (Convertino *et al.* 1993).

In addition to the contribution of reduced plasma volume to the impact of vascular hypovolemia on the reduction in  $\dot{V}O_{2peak}$  after exposure to HDT, the fit subjects demonstrated a significant loss in red blood cell volume that was not observed in the unfit group. Decreased red blood cell volume might have contributed to the greater reduction in  $\dot{V}O_{2peak}$  observed in the fit subjects by lowering oxygen carrying capacity of the blood. Other investigators have verified that red cell mass can decrease from 5 to 25% during prolonged bedrest (Convertino 1995, Branch *et al.* 1997) or spaceflight (Fischer *et al.* 1967, Johnson *et al.* 1975). The findings of the present study might suggest that the stimulus for red blood cell production/degradation may be more sensitive in individuals with high aerobic capacity. However, the relationship between changes in red cell mass and changes in  $\dot{V}O_{2peak}$  is tenuous because significant reductions in  $\dot{V}O_{2peak}$  following bedrest have occurred without change in red cell mass (Convertino *et al.* 1986, Williams and Convertino 1988, Convertino 1995). It therefore appears that the primary effect of microgravity-induced hypovolemia on reducing  $\dot{V}O_{2peak}$  is a result of the contraction of the plasma volume with a lesser contribution from reduced red cell mass.

The major finding of this study was that subjects with relatively high aerobic capacity demonstrated larger reductions in  $\dot{V}O_{2peak}$  and blood volume than individuals with lower aerobic capacity. The data of the present investigation supports the notion that a large  $\dot{V}O_{2peak}$  reserve is associated with greater loss of that reserve during adaptation to microgravity compared

with a more sedentary individual with small  $\dot{V}O_{2peak}$  reserve. This relationship suggests that greater levels of physical activity may be required during spaceflight or rehabilitation following inactivity in aerobically fit individuals to minimize detrimental effects associated with deconditioning. However, these results should not be misinterpreted to suggest that it is detrimental to select astronauts who are aerobically fit. On the contrary, fit individuals remain at a physical and physiological advantage because their aerobic power remains greater than their unfit counterparts after adaptation to microgravity and this represents greater reserve to meet a challenge to perform a given absolute work task.

The authors thank Drs D.J. Goldwater and H. Sandler for their administrative support to this study, Greg M. Karst, Scott M. Kinzer, and Donna A. Williams for their technical assistance with data collection, Dee O'Hara and her staff in the NASA-Ames Research Centre Human Research Facility for their valuable contributions to the organization and supervision of the study, and the test subjects for their cooperation.

## REFERENCES

- Branch, J.D., Pate, R.R., Bodary, P.F. & Convertino, V.A. 1997. Red cell volume and [erythropoietin] responses during exposure to simulated microgravity. *Aviat Space Environ Med* in press. *update!*
- Chase, G.A., Grave, C. & Rowell, L.B. 1966. Independence of changes in functional and performance capacities attending prolonged bed rest. *Aerosp Med* 37, 1232–1238.
- Convertino, V.A. 1990. Physiological adaptations to weightlessness: effects on exercise and work performance. *Exerc Sport Sci Rev* 18, 119–165.
- Convertino, V.A. 1991. Blood volume: its adaptation to endurance training. *Med Sci Sports Exerc* 23, 1338–1348.
- Convertino, V.A. 1995. Exercise and adaptation to microgravity environments. In: M.J. Fregly & C.M. Blatteis (eds) *Handbook of Physiology: Environmental Physiology. III. The Gravitational Environment*. Sect. 1, Ch. 36, pp. 815–843. Oxford University Press, New York.
- Convertino, V.A. 1996. Clinical aspects of the control of plasma volume at microgravity and during return to one gravity. *Med Sci Sports Exerc* 28, S45–S52.
- Convertino, V.A. 1997. Cardiovascular consequences of bedrest: effects on maximal oxygen uptake. *Med Sci Sports Exerc* 29, 191–196.
- Convertino, V.A., Engelke, K.A., Ludwig, D.A. & Doerr, D.F. 1996. Restoration of plasma volume after 16 days of head-down tilt induced by a single bout of maximal exercise. *Am. J Physiol (Regulatory Integrative Comp Physiol)* 270, R3–R10.
- Convertino, V.A., Goldwater, D.J. & Sandler, H. 1986. Bedrest-induced peak  $\dot{V}O_2$  reduction associated with age, gender and aerobic capacity. *Aviat Space Environ Med* 57, 17–22.
- Convertino, V.A., Tatro, D.L. & Rogan, R.B. 1993. Renal and cardiovascular responses to water immersion in trained runners and swimmers. *Eur J Appl Physiol* 67, 507–512.

changed  
to  
match  
list!

- Coyle, E.F., Hemmert, M.K. & Coggan, A.R. 1986. Effects of detraining on cardiovascular responses to exercise: role of blood volume. *J Appl Physiol* 60, 95–99.
- Fischer, C.L., Johnson, P.C. & Berry, C.A. 1967. Red blood cell and plasma volume changes in manned spaceflight. *J Am Med Assoc* 200, 579–583.
- Greenleaf, J.E., Bernauer, E.M., Ertl, A.C., Trowbridge, T.S. & Wade, C.E. 1989. Work capacity during 30-days of bed rest with isotonic and isokinetic exercise training. *J Appl Physiol* 67, 1820–1826.
- Greenleaf, J.E., Convertino, V.A. & Mangseth, G.R. 1979. Plasma volume during stress in man: osmolality and red cell volume. *J Appl Physiol* 47, 1031–1038.
- Greenleaf, J.E. & Kozlowski, S. 1982. Reduction in peak oxygen uptake after prolonged bed rest. *Med Sci Sports Exerc* 14, 477–480.
- Johnson, P.C., Kimzey, S.L. & Driscoll, T.B. 1975. Postmission plasma volume and red-cell mass changes in the crews of the first two Skylab missions. *Acta Astronautica* 2, 311–317.
- Levine, B.D. 1993. Regulation of central blood volume and cardiac filling in endurance athletes. The Frank-Starling mechanism as a determinant of orthostatic tolerance. *Med Sci Sports Exerc* 25, 727–732.
- Levine, B.D., Lane, L.D., Watenpugh, D.E., Gaffney, F.A., Buckey, J.C. & Blomqvist, C.G. 1996. Maximal exercise performance after adaptation to microgravity. *J Appl Physiol* 81, 686–694.
- Saltin, B., Blomqvist, G., Mitchell, J.H., Johnson, R.L., Wildenthal, K. & Chapman, C.B. 1968. Response to exercise after bed rest and after training. *Circulation* 38 (Suppl 7), 1–78.
- Saltin, B., Hartley, L.H., Kilbom, A. & Astrand, I. 1969. Physical training in sedentary middle-aged and older men. II. Oxygen uptake, heart rate, and blood lactate concentration at submaximal and maximal exercise. *Scand. J Clin Lab Invest* 24, 323–334.
- Stegemann, J., Von Framing, H.-D. & Schiefeling, M. 1969. The effect of a six-hour immersion in thermodifferent water on circulatory control and work capacity in trained and untrained subjects. *Pflugers Arch* 312, 129–138.
- Taylor, H.L., Henschel, A., Brozek, J. & Keys, A. 1949. Effects of bed rest on cardiovascular function and work performance. *J Appl Physiol* 2, 223–239.
- Williams, D.A. & Convertino, V.A. 1988. Circulating lactate and FFA during exercise: Effect of reduction in plasma volume following simulated microgravity. *Aviat Space Environ Med* 59, 1042–1046.